

METHODS OF TREATING VASCULAR DISEASE ASSOCIATED WITH CYSTATIN C DEFICIENCY

TECHNICAL FIELD OF THE INVENTION

5 This invention relates generally to methods of treating vascular diseases.

BACKGROUND OF THE INVENTION

Atherosclerosis and abdominal aortic aneurysm (AAA) are inflammatory diseases that involve extensive extracellular matrix degradation and vascular wall remodeling. Cardiovascular events correlate with the presence of inflammation, and atheroma tend to rupture at sites of matrix remodeling (Davies *et al.*, 94 Circulation 213-22 (1996); Ghorpade *et al.*, 800 Ann. N.Y. Acad. Sci. 138-50 (1996)).

However, the mechanisms of degradation of the arterial extracellular matrix in these various contexts remain unclear. In particular, cysteine proteases have previously received little consideration in this mechanism, even though macrophages and smooth muscle cells with greatly expanded lysosomal compartments figure prominently in the histopathology of atherosclerotic plaques. Several cysteine endoproteases (cathepsin K, cathepsin L, and cathepsin S, have potent elastolytic activity. Although elastolytic and collagenolytic cysteine proteases may participate in vascular wall remodeling, cathepsin K has only been observed in bone, and only one study has demonstrated cathepsin K mRNA and protein expression in human breast carcinoma (Littlewood-Evans *et al.*, 57 Cancer Res. 5386-390 (1997)).

Although cathepsins are thought to generally reside in and function optimally within acidic lysosomes, cathepsins also function extracellularly at or near the cell surface. Monocyte-derived human macrophages become markedly elastolytic during *in vitro* culture and predominantly use cysteine proteases to degrade extracellular elastin. In the presence of pro-inflammatory cytokines found in atheroma, cultured smooth muscle cells secrete active cathepsin S capable of degrading extracellular elastin (Sukhova *et al.*,

102 J. Clin. Invest. 576-83 (1998)). Changes in the concentration of cathepsins B and C localized to aortic aneurysm have recently been described (Gacko *et al.*, 50(2) Pol. J. Pathol. 83-86 (1999)), as has increased expression of cathepsin B in the advancing edge of colorectal tumors (Hirai *et al.*, 30(6) Hum. Pathol. 680-86 (1999)).

5 By regulating protease activities, protease inhibitors also play a pivotal role in tissue remodeling (*see*, Werb *et al.*, 857 Ann. N.Y. Acad. Sci. 110-18 (1998)). The most abundant extracellular inhibitor of cysteine proteases is cystatin C (*see*, Barrett *et al.*, 120 Biochem. Biophys. Res. Commun. 631-36 (1984)). Cystatin C is expressed in virtually all organs of the body, and is present in high concentration in biological fluids. However, 10 aside from a rare mutation in cystatin C that leads to its precipitation as amyloid in cerebral blood vessels and causes cerebral hemorrhage, no evidence has thus far implicated cystatin C in disease.

While cystatin C preparations have been described for the treatment of viral disease (United States patent 5,432,264), protection of the nervous system from papain 15 (Hiltke *et al.*, 78(8) J. Dent. Res. 1401-409 (1999)), diagnosis of amyloidosis (United States patent 5,270,165), and identification of therapeutic agents to prevent neuronal cell death (United States patent 5,866,318), the role of cystatin C in vascular disease remains unclear. Until now, it has been unknown whether cystatin C expression actually changes *in situ* in diseases in which inflammation is prominent, and how changes in cystatin C 20 levels might be effected. Thus, there remains a need in the art for the elucidation of the role cystatin C plays in vascular disease, including atherosclerosis, and for the development of therapeutics to prevent and treat vascular elastic laminae breakdown.

SUMMARY OF THE INVENTION

25 The invention is directed to methods of preventing and treating vascular diseases, inflammatory disorders, and vascular tumors by inhibiting cysteine proteases. Specifically, the invention is directed to the inhibition of two cysteine proteases active at

sites of vascular injury, cathepsin S and cathepsin K, by cystatin C, but also includes inhibition of other elastolytic proteases, such as cathepsin L.

The pathogenesis of atherosclerosis and abdominal aortic aneurysm involves elastic laminae breakdown. Elastolytic cysteine proteases are overexpressed at sites of arterial elastin damage, but whether endogenous local inhibitors counterbalance these proteases has previously been unknown. While the cysteine protease inhibitor cystatin C is normally expressed in vascular wall smooth muscle cells, cystatin C is severely reduced in both atherosclerotic and aneurysmal aortic lesions. Increased abdominal aortic diameter correlates inversely with serum cystatin C levels. *In vitro*, cytokine-stimulated vascular smooth muscle cells secrete elastolytic cathepsin S and cathepsin K. The elastolytic activity of these enzymes is blocked by as little as 10 ng/ml of recombinant cystatin C or by transforming growth factor- β_1 (TGF- β_1)-induced cystatin C secretion.

The invention provides methods of preventing the development of and treating vascular disease in a subject by administering an inhibitor of a cysteine protease. In one aspect the subject has not been diagnosed with a vascular disease. Examples of vascular diseases include atherosclerosis or aneurysmal aortic lesions. Cysteine proteases include cystatins, *e.g.* cystatin C polypeptide.

In another aspect the invention provides methods of diagnosing a subject at risk of or with vascular disease. A subject at risk is a subject who for example has a family history of vascular disease or has been diagnosed as having a predisposition to vascular disease (*e.g.*, familial hypercholesterolemia or hereditary angiodema) In addition a subject at risk also may be an individual who demonstrates behavioral or environmental risk factors associated with vascular disease. Such factors include for example, gender, advancing age, cigarette smoking, high blood pressure, diabetes, obesity, lack of physical activity, abnormal blood cholesterol homocysteine levels or decreased serum cystatin C levels as compared to a normal healthy adult value.. The more risk factors a person has, the greater the likelihood of developing vascular disease. Several of these risk factors are

interrelated. Obesity, lack of exercise, and cigarette smoking can raise blood pressure and adversely influence blood cholesterol levels.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a scatter graph illustrating the correlation between aortic diameter and serum cystatin C levels. A cohort of outpatients (n =122) underwent ultrasound measurements of their abdominal aorta and the diameters were correlated with serum cystatin C levels in blood drawn at the time of the outpatient visit. A similar correlation ($p < .04$) was observed with indexed aortic diameter (diameter/body surface area) or with aortic diameter as a function of cystatin C corrected for serum creatinine, a known determinant of cystatin C levels.

FIG. 2 is a bar graph showing that smooth muscle cells elastase activity is inhibited by recombinant cystatin C. Interferon- γ stimulated smooth muscle cells were co-cultured in serum-free medium with insoluble elastin in the presence of increasing concentrations of cystatin C as indicated in the panel. After 72 hrs, media were collected and soluble radioactivity measured as an index of elastin degradation.

FIG. 3 is a graph showing that TGF- β_1 blocks smooth muscle cells elastase activity. Interferon-stimulated smooth muscle cells were co-cultured with elastin as above in the presence of increasing amounts of TGF- β_1 and elastin degradation quantified after 72 hrs.

DETAILED DESCRIPTION OF THE INVENTION

Introduction. Protease inhibitors play a pivotal role in tissue remodeling by regulating protease activity (Werb *et al.*, 857 Ann. N.Y. Acad. Sci. 110-18 (1998)). Among such regulators, the most abundant extracellular inhibitor of cysteine proteases is cystatin C, a 13 kDa protein constitutively secreted shortly after its synthesis (Merz *et al.*, 173 J. Cell Physiol. 423-32 (1997)). Cystatin C belongs to the Type 2 cystatin gene family which cluster, together with other members including cystatins D, 5, SA, and SN,

on chromosome 2 (Abrahamson *et al.*, 82 Hum. Genet. 223-26 (1989)). Cystatins D, S, and SA are primarily expressed in salivary glands whereas cystatin C is expressed in virtually all organs of the body. Due to its high concentration in biological fluids, cystatin C is probably one of the most important extracellular inhibitors of cysteine proteases.

5 Despite the prevalence of cystatin C, no evidence had previously elucidated its role in disease, aside from a rare mutation in cystatin C that leads to its precipitation as amyloid in cerebral blood vessels and causes cerebral hemorrhage. It has additionally been shown that, *in vitro*, alveolar macrophages from cigarette smokers or monocytes stimulated by interferon- γ secrete less cystatin C than unstimulated

10 monocyte/macrophages, raising the possibility of reduced cystatin C levels at sites of alveolar inflammation (Chapman *et al.*, 141 Am. Rev. Respir. Dis. 698-705 (1990); Warfel *et al.*, 166 J. Exp. Med. 1912-917 (1987)). The imbalance of proteolytic and anti-proteolytic enzyme levels may result in lung destruction (Snider, 146 Am. Rev. Respir. Dis. 1615-622 (1992); Tetley, 48 Thorax 560-65 (1993)), tumor progression

15 (DeClerck *et al.*, Adv. Exp. Med. Biol. 425:89-97 (1997); Calkins *et al.*, 376 Biol. Chem. Hoppe Seyler 71-80 (1995)), and possibly vascular wall remodeling (Allaire *et al.*, 102 J. Clin. Invest. 1413-20 (1998); Fabunmi *et al.*, 83 Cir. Res. 270-78 (1998); Knox *et al.*, 95 Circulation 205-12 (1997); Thompson *et al.*, 96 J. Clin. Invest. 318-26 (1995); McMillan *et al.*, 15 Arteriocler. Thromb. Vasc. Biol. 1139-44 (1995)).

20 We have discovered that cystatin C is deficient in diseased arteries. As opposed to normal vascular smooth muscle cells, which highly express cystatin C, immunohistochemical analyses of samples from atherosclerotic plaques revealed virtually no cystatin C antigen within plaques. The severe depression of cystatin C antigen levels within smooth muscle cells of atherosclerotic plaques implies a marked change in the

25 functional properties of plaque smooth muscle cells. Normal medial smooth muscle cells express abundant cystatin C. Such changes in the phenotypic properties of these smooth muscle cells are consistent with other evidence that migrating vascular smooth muscle

cells have protease and protease inhibitor profiles distinctly different from resident cells of the vessel wall. Smooth muscle cells in atheromatous arteries have augmented matrix metalloproteinase levels (Schonbeck *et al.*, 81 Circ. Res. 448-54 (1997); Galis *et al.*, 748 Ann. N.Y. Acad. Sci. 501-7 (1995)). Macrophages within atherosclerotic and aneurysmal tissues also exhibit increased expression of matrix metalloproteinases (Libby *et al.*, 7 Curr. Opin. Lipidol. 330-35 (1996); Pearce *et al.*, 800 Ann. N.Y. Acad. Sci. 175-85 (1996)).

The marked suppression of cystatin C concurrent with augmented expression of cysteine proteases in atherosclerosis and abdominal aneurysms represents the first acquired cysteine protease inhibitor deficiency in human disease. In contrast to results reported here for cystatin C, tissue inhibitors of matrix metalloproteinases (TIMP) are either unchanged or overexpressed in both atherosclerotic lesions, and abdominal aortic aneurysm. Inflammatory cytokines and growth factors increase the expression of TIMP-1 and TIMP-3 in atherosclerotic lesion-related macrophages and vascular smooth muscle cells (Fabunmi *et al.*, 83 Cir. Res. 270-78 (1998)).

Surprisingly, we have found that TGF- β_1 is a potent inducer of cystatin C secretion by vascular smooth muscle cells. Low TGF- β_1 activity explains low levels of lesional cystatin C, and the inverse correlation with aortic dilatation, indicating an additional mechanism by which TGF- β_1 deficiency contributes to the pathobiology of vascular wall remodeling and atherogenesis. Since IFN- γ -stimulated smooth muscle cells secrete elastolytic cathepsin S, and IFN- γ is present within diseased arteries (Hansson *et al.*, 135 Am. J. Path. 169-75 (1989); Szekanecz *et al.*, 42 Agents Actions 159-62 (1994)), secreted cysteine proteases overwhelm residual pericellular cystatin C. The restoration of a physiological balance between the elastolytic cathepsins and inhibitors of these proteases presents a novel therapeutic strategy, especially in aneurysmal disease, where matrix breakdown is dramatic.

In yet another embodiment, the invention provides a method of treating vascular disease in a subject by administering a therapeutically effective amount of TGF- β_1 to the subject. Similarly, the invention also provides a method of preventing vascular disease or blood vessel elastic laminae breakdown in a subject by administering a prophylactically effective amount of TGF- β_1 to the subject. The administration of TGF- β_1 induces secretion of cystatin C. In a particular embodiment, the elastic laminae and collagen breakdown results from inflammation associated with vascular injury.

Cystatin C compositions. In one embodiment, the invention provides a method of treating vascular disease in a subject by administering a therapeutically effective amount of a cysteine proteinase inhibitor to the subject. The invention also provides a method of preventing vascular disease or blood vessel elastic laminae breakdown in a subject by administering a prophylactically effective amount of a cysteine proteinase inhibitor to the subject. The invention also provides a method of preventing elastic laminae and collagen breakdown resulting from inflammation associated with vascular injury by administering a prophylactically effective amount of a cysteine proteinase inhibitor to the subject.

In various embodiments, the cysteine protease inhibitor can be a cathepsin inhibitor. The cathepsin inhibitor may be structurally similar to cystatin C. By structurally similar to cystatin C is meant that the inhibitor contains a cystatin C active site region. (see for example, Hiltke *et al.*, 78(8) J. Dent. Res. 1401-9 (1999) and Mason *et al.*, 330 (Pt. 2) Biochem J. 833-8 (1998).

For example, the cystatin C inhibitor includes the sequence:

[GSTEQKRV]-Q-[LIVT]-[VAF]-[SAGQ]-G-x-[LIVMNK]-x(2)-[LIVMFY]-x-[LIVMFYA]-[DENQKRHSIV] (SEQ ID NO:3)

Alternatively, the cystatin C inhibitor includes the amino acids comprising the amino terminal β -hairpin loop 1 and β -hairpin loop 2 of cystatin C. Hiltke *et al.*, 78(8) J. Dent. Res. 1401-9 (1999).

Mason *et al.* found that amino acid substitutions in the N-terminal segment of cystatin C create selective protein inhibitors of lysosomal cysteine proteinases, using site-directed mutagenesis to alter the specificity of human cystatin C. Olsson *et al.*, 1432(1) Biochim Biophys Acta 73-81 (1999) investigated the affinity and kinetics of inhibition of cysteine proteinases by intact recombinant bovine cystatin C. Furthermore, Nycander *et al.*, 422(1) FEBS Lett. 61-4 found that a two-step mechanism of inhibition of cathepsin B by cystatin C due to displacement of the proteinase occluding loop, using stopped-flow kinetics.

The polypeptide sequence for human cystatin C (EMBL locus HSCYSTC1, Gen Bank Accession No.CAA43856) is
MAGPLRAPLLLLAILAVALAVSPAAGSSPGKPPRLVGGPMDASVEEEGVRRALD
FAVGEYNKASNDMYHSRALQVVRARKQIVAGVNYFLDVELGRTTCTKTQPNLD
NCPFHDQPHLKRKAFCSFQIYAVPWQGTMTLSKSTCQDA (SEQ ID NO:1).

Compositions of cystatin C may be prepared as described in United States patents 5,164,295, 5,270,165; and 5,432,264, incorporated herein by reference.

The cysteine protease inhibitor is cystatin C, or functional equivalents or variants of cystatin C, including recombinant forms. The term "cystatin C" means native or recombinant cystatin C, and functional mutants, variants, or truncated forms thereof.

The polypeptide or nucleic acids of the invention need not be identical to the amino acid residue sequence of the cystatin C polypeptide, for example, so long as the polypeptides are able to bind to and inactivate relevant cathepsins. The polypeptide of the invention also includes any analog, fragment or chemical derivative of the cystatin C polypeptide, so long as the polypeptide is functionally active. The term "analog" includes a polypeptide having an amino acid residue sequence substantially identical to the cystatin C polypeptide sequence in which one or more amino acid residues have been conservatively substituted with a functionally similar residue. Peptide equivalents can differ from the cystatin C polypeptide polypeptides by the substitution or modification of

side chains or functional groups, without destroying biological function. Modifications can include, for example, additions, deletions, or substitutions of amino acids residues, substitutions with compounds that mimic amino acid structure or functions, and the addition of chemical moieties, such as amino or acetyl groups. An "amino acid residue" is an amino acid formed upon chemical digestion (hydrolysis) of a polypeptide at its peptide linkages. A "fragment" is a polypeptide of the invention having an amino acid residue sequence shorter than 50 amino acids.

The amino acid residues described herein are preferably in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the polypeptides inhibit cathepsins. This definition includes, unless otherwise specifically indicated, chemically modified amino acids, including amino acid analogs (such as penicillamine, 3-mercapto-D-valine), naturally occurring non-proteogenic amino acids (such as norleucine), and chemically synthesized compounds that have properties known in the art to be characteristic of an amino acid. The term "proteogenic" indicates that the amino acid can be incorporated into a protein in a cell through well-known metabolic pathways.

The term "conservative substitution" also includes the use of a chemically derivatized residue in place of a non-derivatized residue provided that such polypeptide displays the requisite activity. "Chemical derivative" refers to a polypeptide of the invention having one or more residues chemically derivatized by reaction of a functional side group. Such derivatized molecules include for example, those molecules in which free amino groups have been derivatized to form amine hydrochlorides, p-toluene sulfonyl groups, carbobenzoxy groups, t-butyloxycarbonyl groups, chloroacetyl groups or formyl groups. Free carboxyl groups may be derivatized to form salts, methyl and ethyl esters or other types of esters or hydrazides. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. Also included as chemical derivatives are those peptides that contain one or more naturally occurring amino acid derivatives of the

twenty standard amino acids. For examples, 4-hydroxyproline may be substituted for proline; 5-hydroxylysine may be substituted for lysine; 3-methylhistidine may be substituted for histidine; homoserine may be substituted for serine; and ornithine may be substituted for lysine.

5 Some useful modifications are designed to increase the stability of the polypeptide in solution, and therefore serve to prolong half life of the polypeptides in solutions, particularly biological fluids such as blood, plasma or serum, by blocking proteolytic activity in the blood. A polypeptide can have a stabilizing group at one or both termini. Typical stabilizing groups include amido, acetyl, benzyl, phenyl, tosyl, alkoxycarbonyl, 10 alkyl carbonyl, benzyloxycarbonyl and the like end group modifications. Additional modifications include using a "L" amino acid in place of a "D" amino acid at the termini, cyclization of the polypeptide, and amide rather than amino or carboxy termini to inhibit exopeptidase activity.

 A polypeptide of the invention may or may not be glycosylated. The polypeptides 15 are not glycosylated, for example, when are produced directly by peptide synthesis techniques or are produced in a prokaryotic cell transformed with a recombinant polynucleotide of the invention. Eukaryotically-produced peptide molecules are typically glycosylated.

TGF- β compositions. The term "TGF- β_1 " means native or recombinant TGF- β_1 , 20 and functional mutants, variants, or truncated forms thereof. The sequence of human transforming growth factor-beta, (locus HUMTGFBC accession M60315.1) is
MPGLGRRQWLCWWGLLCSCCGPPPLRPPLPAAAAAAGGQLLDGGSPGR
TEQPPSPQSSSGFLYRRLKTQEKREMQKEILSVLGLPHRPRPLHGLQQPQPPALR
QQEEQQQQQLPRGEPPPGR LKSAPLFMLDLYNALSADNDEDGASEGERQQSW
25 PHEAASSSQRRQPPPGAAHPLNRKSLLAPGSGSGGASPLTSAQDSAFLNDADMV
MSFVNLVEYDKEFSRQRHHKEFKFNLSQIPEGEVVTAAEFRIYKDCVMGSFKN
QTFLISIQVLQEHQHRDSDLFLDTRVWVWASEEGWLEFDITATSNLWVVTPQHN

MGLQLSVVTRDGVHVHPRAAGLVGRDGPYDKQPFMVAFFKVSEVHVRTTRSAS
SRRRQQSRNRSTQSQDVARVSSASDYNSELKTACRKHELTVSFQDLGWQDWII
APKGYAANYCDGECFPLNAHMNATNHAIVQTLVHLMNPEYVPKPCCAPTKLN
AISVLYFDDNSNVILKKYRNMVVRACGCH (SEQ ID NO:2).

5 *How to make the cystatin C or TGF- β polypeptides of the invention.* The polypeptides of the invention can be produced by well known chemical procedures, such as solution or solid-phase peptide synthesis, or semi-synthesis in solution beginning with protein fragments coupled through conventional solution methods, as described by Dugas & Penney, *Bioorganic Chemistry*, 54-92 (Springer-Verlag, New York, 1981).

10 Alternatively, a polypeptide of the invention can be synthesized by using well known methods including recombinant methods and chemical synthesis.

 A polypeptide of the invention can be chemically synthesized, for example, by the solid phase peptide synthesis of Merrifield *et al.*, 85J. Am. Chem. Soc. 2149 (1964). Alternatively, a polypeptide of the invention can be synthesized using standard solution
15 methods (*see, for example*, Bodanszky, *Principles of Peptide Synthesis* (Springer-Verlag, 1984)). Newly synthesized peptides can be purified, for example, by high performance liquid chromatography (HPLC), and can be characterized using, for example, mass spectrometry or amino acid sequence analysis.

 The polypeptides of the invention are particularly useful when they are
20 maintained in a constrained secondary conformation. The terms "constrained secondary structure," "stabilized" and "conformationally stabilized" indicate that the peptide bonds comprising the peptide are not able to rotate freely but instead are maintained in a relatively fixed structure. A method for constraining the secondary structure of a newly synthesized linear peptide is to cyclize the peptide using any of various methods well
25 known in the art. For example, a cyclized polypeptide of the invention can be prepared by forming a peptide bond between non-adjacent amino acid residues as described, for example, by Schiller *et al.*, 25 Int. J. Pent. Prot. Res. 171 (1985). Peptides can be

synthesized on the Merrifield resin by assembling the linear peptide chain using N α -Fmoc-amino acids and Boc and tertiary-butyl proteins. Following the release of the peptide from the resin, a peptide bond can be formed between the amino and carboxy termini.

5 A newly synthesized linear peptide can also be cyclized by the formation of a bond between reactive amino acid side chains. For example, a peptide containing a cysteine-pair can be synthesized, with a disulfide bridge can be formed by oxidizing a dilute aqueous solution of the peptide with K₃Fe(CN)₆. Alternatively, a lactam such as an ,-(glutamyl)-lysine bond can be formed between lysine and glutamic acid residues, a
10 lysinonorleucine bond can be formed between lysine and leucine residues or a dityrosine bond can be formed between two tyrosine residues. Cyclic peptides can be constructed to contain, for example, four lysine residues, which can form the heterocyclic structure of desmosine (*see, for example, Devlin, Textbook of Biochemistry 3rd Edition (1992).* Methods for forming these and other bonds are well known in the art and are based on
15 well known rules of chemical reactivity (Morrison & Boyd, *Organic Chemistry, 6th Edition* (Prentice Hall, 1992)).

Recombinant methods of producing a polypeptide of the invention through the introduction of a vector including a polypeptide encoding the peptide into a suitable host cell is well known in the art, such as is described in Sambrook *et al., Molecular Cloning: A Laboratory Manual, 2nd Edition* (Cold Spring Harbor Press, 1989). The term
20 "recombinant" refers to the molecular biological technology for combining polynucleotides to produce useful biological products, and to the polynucleotides and peptides produced by this technology. The polynucleotide can be a recombinant construct (such as a vector or plasmid) which contains the polynucleotide encoding the peptide or
25 fusion protein under the operative control of polynucleotides encoding regulatory elements such as promoters, termination signals, and the like. "Operatively linked" refers to a juxtaposition wherein the components so described are in a relationship permitting

09697613-102600
099201-ET926960

them to function in their intended manner. A control sequence operatively linked to a coding sequence is ligated such that expression of the coding sequence is achieved under conditions compatible with the control sequences. "Control sequence" refers to polynucleotide sequences which are necessary to effect the expression of coding and non-coding sequences to which they are ligated. Control sequences generally include promoter, ribosomal binding site, and transcription termination sequence. Determinations of the sequences for the polynucleotide coding region that codes for the peptides described herein can be performed using commercially available computer programs, such as DNA Strider and Wisconsin GCG. Owing to the natural degeneracy of the genetic code, the skilled artisan will recognize that a sizable yet definite number of DNA sequences can be constructed which encode the claimed peptides (*see, Watson et al., Molecular Biology of the Gene*, 436-437 (The Benjamin/Cummings Publishing Co., 1987)).

In addition, "control sequences" refers to sequences which control the processing of the peptide encoded within the coding sequence; these can include, but are not limited to sequences controlling secretion, protease cleavage, and glycosylation of the peptide. The term "control sequences" is intended to include, at a minimum, components whose presence can influence expression, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences. A "coding sequence" is a polynucleotide sequence that is transcribed and translated into a polypeptide. Two coding polynucleotides are "operably linked" if the linkage results in a continuously translatable sequence without alteration or interruption of the triplet reading frame. A polynucleotide is operably linked to a gene expression element if the linkage results in the proper function of that gene expression element to result in expression of the coding sequence. "Transformation" is the insertion of an exogenous polynucleotide (*i.e.*, a "transgene") into a host cell. The exogenous polynucleotide is integrated within the host genome. A polynucleotide is "capable of expressing" a peptide if it contains

nucleotide sequences which contain transcriptional and translational regulatory information and such sequences are "operably linked" to polynucleotide which encode the peptide. A polynucleotide that encodes a peptide coding region can be then amplified, for example, by preparation in a bacterial or viral vector, according to conventional methods, for example, described in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Edition (Cold Spring Harbor Press, 1989). Expression vehicles include plasmids or other vectors.

The polynucleotide encoding the polypeptide of the invention can be prepared by chemical synthesis methods or by recombinant techniques. The polypeptides can be prepared conventionally by chemical synthesis techniques, such as described by Merrifield, 85 J. Amer. Chem. Soc. 2149-2154 (1963) (*see*, Stemmer *et al.*, 164 Gene 49 (1995)). Synthetic genes, the *in vitro* or *in vivo* transcription and translation of which will result in the production of the protein can be constructed by techniques well known in the art (*see* Brown *et al.*, 68 Methods in Enzymology 109-151 (1979)). The coding polynucleotide can be generated using conventional DNA synthesizing apparatus such as the Applied Biosystems Model 380A or 380B DNA synthesizers (commercially available from Applied Biosystems, Inc., 850 Lincoln Center Drive, Foster City, Calif. 94404).

Alternatively, systems for cloning and expressing polypeptide of the invention include various microorganisms and cells that are well known in recombinant technology. These include, for example, various strains of *E. coli*, *Bacillus*, *Streptomyces*, and *Saccharomyces*, as well as mammalian, yeast and insect cells. Suitable vectors are known and available from private and public laboratories and depositories and from commercial vendors. *See*, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Edition (Cold Spring Harbor Press, 1989). *See also*, PCT International patent application WO 94/01139). These vectors permit infection of patient's cells and expression of the synthetic gene sequence *in vivo* or expression of it as a peptide or fusion protein *in vitro*.

Polynucleotide gene expression elements useful for the expression of cDNA encoding peptides include, but are not limited to (a) viral transcription promoters and their enhancer elements, such as the SV40 early promoter, Rous sarcoma virus LTR, and Moloney murine leukemia virus LTR; (b) splice regions and polyadenylation sites such as those derived from the SV40 late region; and (c) polyadenylation sites such as in SV40. Recipient cells capable of expressing the gene product are then transfected. The transfected recipient cells are cultured under conditions that permit expression of the gene products, which are recovered from the culture. Host mammalian cells, such as Chinese Hamster ovary cells (CHO) or COS-1 cells, can be used. These hosts can be used in connection with poxvirus vectors, such as vaccinia or swinepox. Suitable non-pathogenic viruses that can be engineered to carry the synthetic gene into the cells of the host include poxviruses, such as vaccinia, adenovirus, retroviruses and the like. A number of such non-pathogenic viruses are commonly used for human gene therapy, and as carrier for other vaccine agents, and are known and selectable by one of skill in the art. The selection of other suitable host cells and methods for transformation, culture, amplification, screening and product production and purification can be performed by one of skill in the art by reference to known techniques (*see, e.g.*, Gething & Sambrook, 293 Nature 620-625 (1981)).

Another preferred system includes pSecTag2 expression vectors (Invitrogen, San Diego, CA). These 5.2 kb mammalian expression vectors offer features for secretion, purification, and detection of fusion proteins. The vectors carry the secretion signal from the V-J2-C region of the mouse Ig 6-chain for efficient secretion of recombinant proteins from a variety of mammalian cell lines. The vectors also carry a C-terminal tag sequence that allows rapid purification and detection of fusion proteins. Recombinant proteins are fused to six histidine residues to allow one-step purification (which can be removed following purification) and the anti-*myc* epitope for rapid detection.

When produced by conventional recombinant means, peptide can be isolated either from the cellular contents by conventional lysis techniques or from cell medium by conventional methods, such as chromatography (*see, e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Edition* (Cold Spring Harbor Press, 1989)). Methods for recovering an expressed recombinant polypeptide are well known in the art and include fractionation of the protein-containing portion of the culture using well known biochemical techniques. For instance, the methods of gel filtration, gel chromatography, ultrafiltration, electrophoresis, ion exchange, affinity chromatography and the like, such as are known for protein fractionations, can be used to isolate the expressed proteins found in the culture. In addition, immunochemical methods, such as immunoaffinity, immunoadsorption and the like can be performed using well known methods.

Pharmaceutical compositions. The invention provides pharmaceutical compositions comprising a therapeutically effective dose of the polypeptide and nucleic acids of the invention in a pharmaceutically acceptable excipient, for administration to a subject, such as a human patient. The term "therapeutically effective amount" means the dose needed to effectively treat cellular infiltration and attendant cytokine network alterations associated with a variety of inflammatory diseases and injuries. For purposes of the invention, the terms "treat" and "treatment" include preventing, inhibiting, reducing the occurrence of, or ameliorating the physiological effects of the inflammatory condition treated.

The term "pharmaceutically acceptable excipient" includes any solvents, dispersion media, antibacterial and antifungal agents, microcapsules, liposomes, cationic lipid carriers, isotonic and absorption delaying agents and the like which are not incompatible with the active ingredients. The formulation of pharmaceutical compositions is generally known in the art; reference can conveniently be made to *Remington's Pharmaceutical Sciences, 17th Edition* (Mack Publishing Co., Easton, Pa.) Thus, the pharmaceutical compositions can comprise a suitable application medium, such

as a gel, salve, lotion, colloid or powder, aqueous solutions such as physiologically buffered saline or other solvents or vehicles such as glycols, glycerol, oils such as olive oil or injectable organic esters. Pharmaceutical compositions can be prepared as injectables, either as liquid solutions or suspensions. The preparation can also be emulsified.

Physiologically acceptable excipients include, for example, carbohydrates, such as glucose, sucrose or dextrans, antioxidants, such as ascorbic acid or glutathione, chelating agents, low molecular weight proteins or other stabilizers or excipients. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the composition can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents that enhance the effectiveness of the active ingredient. A polypeptide can also be formulated into the pharmaceutical composition as neutralized pharmaceutically acceptable salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide or antibody molecule) that are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, tartaric, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

The pharmaceutical forms suitable for infusion include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The form is fluid to the extent that easy syringability exists. Typical excipients include a solvent or dispersion medium containing, for example, water buffered aqueous solutions (*i.e.*, biocompatible buffers), ethanol, polyols such as glycerol, propylene glycol, polyethylene glycol, suitable mixtures thereof, surfactants, or vegetable oils. Sterilization can be accomplished by any art-recognized technique, including but not

limited to filtration or addition of antibacterial or antifungal agents, for example, paraben, chlorobutanol, phenol, sorbic acid or thimerosal. Further, isotonic agents such as sugars or sodium chloride may be incorporated in the pharmaceutical compositions. Prevention against microorganism contamination can be achieved through the addition of various antibacterial and antifungal agents.

Methods of administration. The cystatin C and cystatin C expression-inducing compositions (*e.g.* TGF- β_1) of the invention may be administered to a subject in any manner known to those of skill in the art. In particularly preferred embodiments, the administration of cystatin C blocks about 50% or greater of IFN- γ -induced cathepsin S activity, and the administration of TGF- β_1 blocks about 75% of IFN- γ -induced cathepsin S activity.

The polypeptides of the invention can be administered in any way that is medically acceptable which may depend on the disease condition or injury being treated. For example the inhibitors of the invention may be administered locally in a solution form at a site of vascular injury through a catheter port or as a coating on the surface of a catheter, *e.g.*, a perfusion catheter or the balloon portion of a catheter used for balloon catheterization. Possible administration routes include injections, by parenteral routes such as intravascular, intravenous, intra-arterial, subcutaneous, intramuscular, intratumor, intraperitoneal, intraventricular, intraepidural or others, as well as oral, nasal, ophthalmic, rectal, topical, or by inhalation. The compositions may also be directly applied to tissue surfaces during surgery. For example the inhibitors can be administered before, during or after angioplasty. In addition, the inhibitors can be administered as a coating on a stent that is implanted at the site of a vascular lesion, *e.g.*, an aneurysm. (*see e.g.*, USSN 5,902,266) Sustained release administration is also specifically included in the invention, by such means as depot injections or erodible implants. Administration can also be intra-articularly, intrarectally, intraperitoneally, intramuscularly, subcutaneously, or by aerosol inhalant in order to prevent particular inflammatory and immune reactions

associated with such areas of the body. Where treatment is systemic due, for example, to the presence in the subject of metastatic tumor cells, the composition can be administered orally or parenterally, such as intravenously, intramuscularly, subcutaneously, intraorbitally, intracapsularly, intraperitoneally or intracisternally.

5 The pharmaceutical compositions are conventionally administered intravenously, as by injection of a unit dose. The term "unit dose" when used in reference to a pharmaceutical composition of the invention refers to physically discrete units suitable as unitary dosages for humans, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect.

10 The quantity to be administered depends on the subject to be treated, capacity of the subject to utilize the active ingredient, and degree of inhibition of receptor-ligand binding desired. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner and are peculiar to each individual. However, suitable dosage ranges are of the order of one to several mg of active ingredient per individual per
15 day and depend on the route of administration. Suitable regimes for initial administration and booster shots are typified by an initial administration followed by repeated doses at one or more hour intervals by a subsequent injection or other administration. Alternatively, continuous intravenous infusion sufficient to maintain therapeutically effective concentrations in the blood are contemplated. For a polypeptide of the
20 invention, therapeutically effective blood concentrations are in the range of about 1 nM to 1 mM, preferably from 50 nM to 100 μ M.

 Whenever, the polypeptides of the invention are used for promoting attachment of cells, such compositions will typically have a higher concentration than those taken internally.

25 *Dosage.* The precise therapeutically effective amount of polypeptide of the invention used in the methods of this invention applied to humans can be determined by the ordinarily skilled artisan with consideration of individual differences in age, weight,

extent of cellular infiltration by inflammatory cells and condition of the patient. The pharmaceutical preparation of the invention should be administered to provide an effective concentration of 1 nM to 1 mM, preferably from 50 nM to 100 μ M. In preferred embodiments, the therapeutically and prophylactically effective amounts of cystatin C
5 comprise about 10-500 ng/ml, and the therapeutically and prophylactically effective amounts of TGF- β_1 comprise about 1-10 ng/ml. In particularly preferred embodiments, the therapeutically and prophylactically effective amounts of cystatin C and TGF- β_1 comprise about 50 ng/ml and about 1 ng/ml, respectively.

The amount of the inhibitor in the pharmaceutical composition of the present
10 invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of inhibitor with which to treat each individual patient. Initially, the attending physician will administer low doses of inhibitor and observe the patient's response. Larger doses of inhibitor may be administered until the optimal
15 therapeutic effect is obtained for the patient, and at that point the dosage is not generally increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.1 μ g to about 100 mg of inhibitor per kg body weight. Preferred ranges include is a dosages of about 100 μ g to 6 mg, and 20 μ g to 500 μ g per kg of body weight.

The duration of intravenous therapy using the pharmaceutical composition of the
20 present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

The total effective amount of a polypeptide of the invention can be administered
25 to a subject as a single dose, either as a bolus or by infusion over a relatively short period

of time, or can be administered using a fractionated treatment protocol, in which the multiple doses are administered over a more prolonged period of time.

It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required pharmaceutical excipient. A pharmaceutical composition typically contains an amount of at least 0.1 weight % of active ingredient, *i.e.*, a polypeptide or antibody of this invention, per weight of total pharmaceutical composition. A weight % is a ratio by weight of active ingredient to total composition. Thus, for example, 0.1 weight % is 0.1 grams of polypeptide per 100 grams of total composition.

In vitro diagnostics. Patients with progressing atherosclerosis exhibit low circulating cystatin C, even though they may not present gross aortic morphological change. This observation represents a novel means of diagnosing patients at risk of, or with, progressing vascular disease. In one embodiment, the invention provides a method of diagnosing a patient at risk of, or with, vascular injury by detecting low circulating levels of cystatin C in blood serum. In a particularly preferred embodiment the low cystatin C level is about 50% or less of the control cystatin C level (where the ratio is determined using levels defined as the mean \pm 2 standard deviations), and in another particularly preferred embodiment, the vascular injury comprises atherosclerosis (*see*, Randers et al., Clin. Chem. Lab Med., 37(4):389-95 (1999)). For example, the normal patient control cystatin C level in EXAMPLE III is 1.303 ± 1.064 μ g/ml. This detection may be accomplished by any method well known in the art, for example, ELISA, as discussed above.

Therapeutic use. Vascular diseases within the scope of the invention include, but are not limited to, atherosclerosis, myocardial infarction, arterial aneurysms, including

abdominal aortic aneurysm, inflammatory diseases, including giant cell arteritis, vascular tumors, and tumor-induced vascular lesions.

We have discovered a downward skewing of circulating cystatin C levels among subjects with dilated aortas and the severe depletion of cystatin C within atherosclerotic and aneurysmal lesions. This downward skewing indicate a common response to similar local and systemic changes in the cytokine milieu in these settings. Numerous studies have documented evidence of increased local and systemic markers of inflammation, *e.g.* the cytokine IL-6 and the acute phase reactant, C-reactive protein, in atherosclerosis and aneurysmal disease (Szekanecz *et al.*, 42 Agents Actions 159-62 (1994); Ridker *et al.*, 46 J. Investig. Med. 91-95 (1998)). Additionally, both decreased systemic active TGF- β_1 and local resistance to TGF- β_1 activity have been reported in atherosclerosis by Grainger *et al.*, 1 Nature Med. 74-79 (1995); and McCaffrey *et al.*, 100 J. Clin. Invest. 2182-88 (1997).

The method of the invention inhibits the activity of cathepsin K. Cathepsin K is the most potent mammalian elastase yet described (*see*, Bossard *et al.*, 271 J. Biol. Chem. 12517-524 (1996); Bromme *et al.*, 271 J. Biol. Chem. 2126-32 (1996)). Cathepsin K also has collagenolytic activity (Kafienah *et al.*, 331 Biochem J. 727-32 (1998)).

The overexpression of cathepsin S and cathepsin K in atherosclerotic lesions compared to normal arteries has been demonstrated by Shi *et al.*, 357 FEBS Lett. 129-34 (1995). Also, macrophages and smooth muscle cells in atherosclerotic plaques express cathepsins S and K (Sukhova *et al.*, 102 J. Clin. Invest. 576-83 (1998)). In the presence of pro-inflammatory cytokines found in atheroma, cultured vascular smooth muscle cells secrete active cathepsin S capable of degrading extracellular elastin. Monocyte-derived macrophages also degrade elastin through release of active cathepsins (*see*, Reedy *et al.*, 92 Porch. Natl. Acad. Sci. 3849-853 (1995)).

This degradative mechanism is reminiscent of the bone macrophage, osteoclasts. Indeed, in osteoclasts, cathepsin K appears to be the major protease involved in

extracellular matrix metabolism necessary for normal bone growth and remodeling (Bossard *et al.*, 271 J. Biol. Chem. 12517-524 (1996)). Deficiency of this enzyme results in accumulation of bone matrix mostly around, rather than in, osteoclasts, consistent with the primary site of action of cathepsin K in the pericellular space (Gelb *et al.*, 273 Science 1236-38 (1996)). Together, these studies indicate that under some conditions
5 macrophages utilize certain elastolytic cathepsins to remodel extracellular matrix.

Thus, the invention is useful for preventing, inhibiting, or ameliorating inflammatory and immune reactions associated with various injury and disease conditions. The term "subject" means any mammal to which the compositions of the
10 invention may be administered. Subjects specifically intended for treatment with the compositions and methodologies of the invention include humans.

Inflammation is the body's reaction to injury and infection. Three major events are involved in inflammation: (1) increased blood supply to the injured or infected area; (2) increased capillary permeability enabled by retraction of endothelial cells; and (3)
15 migration of leukocytes out of the capillaries and into the surrounding tissue (hereinafter referred to as cellular infiltration; *see*, Roitt *et al.*, *Immunology*, (Grower Medical Publishing, New York, 1989). Inflammatory cells and mediators including leukocytes, cytokines, oxygen radicals, complement and arachidonate metabolite damage capillary endothelium and allow fluid and protein to leak across capillaries. Inflammation also
20 involves the release of inflammatory mediators (inflammatory cytokines, platelet activating factor, complement, leukotrienes, oxygen metabolites, and the like). Inflammatory responses are thus induced both at the original site of trauma and also in the vasculature and remote vascularized sites. Neutrophils, eosinophils, mast cells, lymphocytes and macrophages contribute to the inflammatory response. Neutrophils
25 contribute by generating oxidants or releasing proteases that damage the microvasculature or adjacent tissue. Cell death and tissue damage due to complement and inflammatory

cell mechanisms lead to organ failure or decreased organ function. The activation of mediators by a local injury can also cause a remote injury to highly vascularized organs.

5 The details of one or more embodiments of the invention are set forth in the accompanying description above. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. Other features, objects, and advantages of the invention will be apparent from the description and from the claims. In the specification and the appended claims, the singular forms include plural referents unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. All patents and publications cited in this specification are incorporated by reference.

10 The following EXAMPLES are presented in order to more fully illustrate the preferred embodiments of the invention. These examples should in no way be construed as limiting the scope of the invention, as defined by the appended claims.

EXAMPLE 1
20 IMMUNOSTAINING TO SHOW CYSTATIN C EXPRESSION IN NORMAL AND ATHEROSCLEROTIC HUMAN ARTERIAL TISSUE

This EXAMPLE shows decreased expression of cystatin C in atherosclerotic plaques and abdominal aortic aneurysm (AAA) by the immunostaining of normal vessels and atherosclerotic plaques.

25 Plaques and normal vessels were immunostained with rabbit anti-human cystatin C polyclonal antibodies to examine if the lesions also exhibit altered levels of their endogenous inhibitor cystatin C. Slides for immunostaining were prepared as follows: Serial cryostat sections (6 μ m) of human atherosclerotic plaques from coronary and carotid arteries, aortic aneurysms, as well as nonatherosclerotic arteries (carotids from

00920T ET 92600
09697613 102600

autopsies and aortas from cardiac transplantation donors were fixed in acetone (-20°C, 5 minutes), air dried, and stained by an avidin-biotin-peroxidase method as previously described (Sukhova *et al.*, 102 J. Clin. Invest. 576-83 (1998)). Tissue sections were treated with 0.3% hydrogen peroxide to inhibit endogenous peroxidase activity and incubated with primary antibodies diluted in PBS supplemented with 4% species-appropriate normal serum. The subsequent processing was performed according to the manufacturer's recommendations (Universal Dako LSAB kit, peroxidase; DAKO Co., Carpinteria, California, USA). The reaction was visualized with 3-amino-9-ethylcarbazole (AEC; Sigma Chemical Co., St. Louis, Missouri, USA). Sections were counterstained with Gill's hematoxylin solution (Sigma Chemical Co.). Cell types were identified with monoclonal anti-muscle actin HHF-35 (Enzo Diagnostics Inc., Farmingdale, New York, USA) or monoclonal anti-human CD68 (macrophages).

Normal arteries exhibit abundant cystatin C staining. However, atherosclerotic plaques show little immunoreactive cystatin C. Macrophages and smooth muscle cells in the developing plaques were identified by immunostaining plaques with mouse anti-human CD68 (macrophage-specific) and mouse anti-human α -actin (smooth muscle cells specific) antibodies. As expected, vivid CDG8 staining is found at the lipid core and shoulder areas of the plaque whereas α -actin staining was found mostly in the fibrous cap. The specificity of immunohistochemistry was confirmed using nonimmune rabbit serum.

Surprisingly, the level of cystatin C in atherosclerotic plaques correlates with disease progression. Early stages of plaque development (fatty streaks) show very little immunodetectable cystatin C, whereas smooth muscle cells of underlying media (α -actin positive) demonstrate stronger staining for cystatin C. Abdominal aortic aneurysm displays a similar lack of cystatin C staining, as well as increased cathepsin S immunostaining. Thus, abdominal aortic aneurysm and atherosclerotic plaques contain scant cystatin C.

EXAMPLE 2
IMMUNOBLOTTING TO SHOW CYSTATIN C EXPRESSION IN NORMAL AND
ATHEROSCLEROTIC HUMAN ARTERIAL TISSUE

5 Immunoblotting was performed with normal arteries, atherosclerotic plaques, and aortic aneurysms using rabbit polyclonal antibodies against cystatin C, cathepsin S or cathepsin K. Both atherosclerotic plaques and aortic aneurysms demonstrated markedly decreased cystatin C antigen, whereas both cathepsin S and cathepsin K were increased in aneurysmal aortae.

10 Protein extracts of atherosclerotic artery, aneurysmal tissue, and normal control vessel were examined by immunoblotting with cathepsin and cystatin C antibodies. Western blot analysis was performed as follows: Frozen atherosclerotic plaques, abdominal aortic aneurysm aortas, and normal arteries were pulverized and lysed into buffer containing 1% Triton-X-100, 40 mM sodium acetate, and 1 mM EDTA, pH 5.0.
15 After incubation at 37°C for 1 hr., lysate protein concentrations were determined using Bio-Rad Dc protein assay kit according to the manufacturer (Bio-Rad Dc., Bio-Rad Laboratories, Inc., Hercules, California, USA). Thirty micrograms of protein from each sample was separated on SDS-PAGE, blotted onto nitrocellulose filter, and probed with rabbit anti-human cystatin C (1:1000, DAKO Corporation) or human cathepsin S
20 (1:1000).

Human saphenous vein smooth muscle cells were cultured in 6-well plates in Dulbecco's Modification of Eagle's Medium (DMEM) containing 10% fetal bovine serum (FBS) until confluent. Cells were then cultured in serum free DMEM for 24 hours followed by stimulation with INF- γ (500 units/ml) for additional 24 hours in the presence
25 or absence of 10 ng/ml TGF- β 1. Culture medium was then collected and centrifuged at 16000xg for 10 min. Proteins were precipitated with 20% trichloroacetic acid on ice for 30 min. followed by repeat centrifugation. The protein pellet was briefly washed with 100% ethanol and resuspended in electrophoresis sample buffer. smooth muscle cells were lysed directly into sample buffer after two washes with 1xPBS. Both cell lysates and

proteins from culture medium were separated on SDS-PAGE for immunoblot analysis with antibodies against human cathepsin S and cystatin C as described above. All Western blots were developed with Western Blot Chemiluminescence Reagent (NEN Life Science Products, Inc., Boston, Massachusetts, USA).

5 Similar to the findings with immunostaining, immunoblot analysis with cystatin C antibodies indicates a dramatic decrease of 13 kDa cystatin C antigen in atherosclerotic plaque extracts, as well as abdominal aortic aneurysm extracts, as compared with normal arteries. Cathepsins S and K expression in abdominal aortic aneurysm extracts is increased, consistent with previously reported data for atherosclerotic plaques (Sukhova
10 *et al.*, 102 J. Clin. Invest. 576-83 (1998)).

These findings evidence a disturbed balance between proteases and anti-proteases involving elastolytic and collagenolytic cysteine proteases in these diseased vessel walls.

EXAMPLE 3
15 CORRELATION BETWEEN ABDOMINAL AORTIC ANEURYSM AORTA
DIAMETER AND SERUM CYSTATIN C LEVELS

A cohort of 4 patients referred to an outpatient cardiology clinic for echocardiographic testing underwent measurements of carotid artery intimal to medial
20 thickness (IMT) ratios and aortic diameter. Correlations between these measurements and circulating indices of inflammation in this cohort have been previously reported (Rhode
et al., 18 Arterioscler. Thromb. Vasc. Biol. 1765-70 (1998)). Serum cystatin C levels in these patients were measured using a cystatin C enzyme-linked immunoassay (ELISA). This method has been previously used to quantify cystatin C levels in human gingiva
25 (Skaleric *et al.*, 34 Arch. Oral. Biol. 301-305 (1989); Shimode *et al.*, 22 Stroke 860-66 (1991); Ishiguro *et al.*, 8 Hybridoma 303-13 (1989)).

The patient population was as follows: Outpatients older than 50 years of age, referred to the Noninvasive Cardiac Laboratory of the Brigham and Women's Hospital for a transthoracic echocardiogram, were invited to participate in a study examining

009207 17926950
5 markers of vascular disease (Rhode *et al.*, 18 Arterioscler. Thromb. Vasc. Biol. 1765-70 (1998)). Patients with a clinical history consistent with active infection, systemic inflammatory disease, heart transplant or those taking corticosteroids were excluded. Overall, 122 eligible patients agreed to participate and constituted the study population. 5
10
15
20
25
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
22

No significant correlation between IMT and cystatin C levels was apparent. However, a significant negative correlation of abdominal aortic diameters with serum cystatin C levels was observed (FIG. 1) ($P < .03$, $R = .203$). The correlation remained significant when aortic diameter was corrected for body surface area and when cystatin C was normalized to serum creatinine, glomerular filtration being a known determinant of cystatin C levels (Kyse-Andersen *et al.*, 40 Clin. Chem. 1921-26 (1994)).

All patients with dilated aortas had relatively low circulating cystatin ($0.516 \pm 0.233 \mu\text{g/ml}$ as compared to normal patients $1.303 \pm 1.064 \mu\text{g/ml}$; *see* FIG. 1), although the number of patients with frankly dilated aortas ($> 2.5 \text{ cm}$) is small. As a group these subjects had significantly lower serum cystatin C levels than that of subjects with aortic diameters $< 2.5 \text{ cm}$ ($P < .03$, Mann-Whitney). There is no significant association of cystatin C with age, race, sex, history of smoking, diabetes, myocardial infarction, revascularization, or lipoprotein levels. Hence, cystatin C levels contribute to progression of vascular remodeling, consistent with significantly lower cystatin C levels among subjects with dilated aortas. Additionally, cystatin C exerts a protective (*i.e.* preventative) effect against matrix breakdown (FIG. 1).

EXAMPLE 4 CYSTATIN C EXPRESSION AND REGULATION OF VASCULAR SMOOTH MUSCLE CELLS ELASTOLYTIC ACTIVITY

Prior studies documenting the capacity of macrophages to use cysteine proteases to degrade elastin found macrophage elastolytic activity to be insensitive to the presence of pericellular serum protease inhibitors (Reedy *et al.*, 92 *Proc. Natl. Acad. Sci.* 3849-853 (1995); Chapman *et al.*, 74 *J. Clin. Invest.* 1693-700 (1984)). Given the marked reduction of cystatin C in atherosclerotic and aneurysmal lesions and the clinical correlation in EXAMPLES 1-3, the influence of cystatin C on the elastolytic activity of cultured vascular smooth muscle cells was examined. Vascular smooth muscle cells were stimulated with IFN- γ (500 units/ml) for 24 hours to augment cathepsin S-dependent

elastase activity (Sukhova *et al.*, 102 J. Clin. Invest. 576-83 (1998)). In co-cultures with insoluble elastin, the addition of as little as 10 ng/ml of recombinant cystatin C inhibits elastin degradation (FIG. 2). Higher concentrations of cystatin C block >75% of the IFN- γ -induced activity, confirming that smooth muscle cells elastase activity depends largely on elastolytic cysteine proteases.

The regulation of cystatin C expression in smooth muscle cells was further tested using a cytokine previously reported to induce other classes of protease inhibitors, TGF- β (Buisson *et al.*, 12 FASEB J. 1683-91 (1998); Cao *et al.*, 36 Invest. Ophthalmol. Vis. Sci. 1411-19 (1995); Su *et al.*, 70 J. Cell. Biochem. 517-27 (1998)).

Total RNAs were isolated from control and IFN- γ -stimulated smooth muscle cells for Northern blot analysis. Northern blot analysis was performed as follows: After INF- γ and TGF- β 1 treatments, 1.5×10^7 smooth muscle cells were lysed in 5 ml of guanidium isothiocyanate (GTC) lysis buffer for RNA preparation as described (Sukhova *et al.*, 102 J. Clin. Invest. 576-83 (1998)). Twenty micrograms of total RNA from smooth muscle cells were separated on 1.2% agarose gel and blotted on Zeta-Probe GT blotting membrane (Bio-Rad Laboratories, Inc.). A human cystatin C probe was generated by polymerase chain reaction using cystatin C cDNA. The 500 bp cystatin C cDNA fragment and cathepsin S cDNA probe (Sukhova *et al.*, 102 J. Clin. Invest. 576-83 (1998)) were labeled with [32 P]-dCTP and purified with NENSORB 20 column (NEN Life Science Products, Inc.) before hybridizing as previously described (Sukhova *et al.*, 102 J. Clin. Invest. 576-83 (1998)).

The elastase assay was performed as follows: Smooth muscle cells were cultured on 24-well plates in 10% FBS DMEM until 100% confluence (5×10^5 cells/well). Cells were then cultured in serum free DMEM for 24 hr before stimulation with INF- γ (500 Units/ml) and/or TGF- β 1 (1-25 ng/ml). Both stimulated and non-stimulated smooth muscle cells were incubated with 300 μ g [3 H]-elastin (bovine ligament elastin, 1300 cpm/ μ g, Elastin Products Inc.) as previously described (Sukhova *et al.*, 102 J. Clin. Invest.

576-83 (1998)) for 3 days at 37°C. The inhibitory effect of cystatin C was examined by incubating smooth muscle cells with recombinant cystatin C (0-500 ng/ml) or rabbit anti-human cystatin C IgG (100 µg/ml, Cortex Biochem, San Leandro, California, USA). Culture media were then collected by centrifugation at 16,000xg for 15 min. The digested elastin radioactivity was counted using 200 µl of medium and the data presented as µg elastin degraded per 10⁶ cells in 24 hours.

As expected, cathepsin S mRNA is markedly increased in stimulated cells as compared with controls.

Surprisingly, the level of cystatin C mRNA was unchanged, consistent with the constitutive nature of the cystatin C promoter (Olafsson, 55 Scand. J. Clin. Lab. Invest. 597-607 (1995)).

TGF-β₁ (5 ng/ml) does not change the level of cathepsin S mRNA or protein, or cystatin C mRNA. However, TGF-β₁ markedly increases the level of secreted cystatin C. Cystatin C biosynthesis increases as little as one hour after exposure of smooth muscle cells to TGF-β₁. Consistent with this observation, as little as 1 ng/ml of recombinant TGF-β₁ almost completely blocks the elastase activity of IFN-γ-stimulated smooth muscle cells (*see*, FIG. 3). Thus, cytokine-induced smooth muscle cells elastase activity depends on pericellular cystatin C levels and TGF-β₁ can induce cystatin C secretion.

In summary, TGF-β₁ stimulates smooth muscle cells cystatin C secretion, while IFN-γ induces cathepsin S mRNA and secretion of active cathepsin S, but has no effect on cystatin C secretion.

EXAMPLE 5 IN VIVO SUPPRESSION OF CATHEPSIN ACTIVITY BY CYSTATIN C

The ability of cystatin C to inhibit the activity of cathepsin S and K *in vivo* is studied using well known atherosclerosis animal models (*see*,; Purcell-Huynh *et al.*, 95 J. Clin. Invest. 2246-257 (1995); Fruebis *et. al.*, 94 J. Clin. Invest. 392-98 (1994); van Ree

et al., 305 *Biochem J.* 905-11 (1995); Masucci *et al.*, 275 *Science* 391-394 (1997); Chowdhury *et al.*, 254 *Science* 1802-805 (1991); Tangirala *et al.*, 36 *J. Lipid Res.* 2320-328 (1995); Ishibashi *et al.* 93 *J. Clin. Invest.* 1885-893 (1994)).

In particular, the LDLR^{-/-} mouse model of Litchman *et al.*, 154 *FASEB J.* 896 (1997) is used, in which atherosclerotic lesion formation in these mice are initiated in a defined fashion by dietary manipulation. Male mice are used, because female mice are not as responsive to diet-induced atherogenesis. Systemic administration of the cathepsins S and K inhibitor cystatin C during the period of atherogenesis are carried out, and animals are sacrificed, and their hearts and aortas collected. Aortae from both cystatin C treated and untreated mice are analyzed with immunohistochemistry to quantify lesional development as above in EXAMPLE 1.

Also, the LDLR^{-/-} mice are manipulated with cathepsin S and cathepsin K transgenes. Mouse cathepsin S and K cDNAs driven by the mouse macrophage scavenger receptor promoter are used to generate a transgenic mouse. Overexpression of cathepsins S and K in LDLR^{-/-} mouse is performed as follows: Both mouse cathepsin S and cathepsin K cDNAs are subcloned, respectively, into the 3'-end of mouse macrophage scavenger receptor, a promoter which has been shown to drive gene expression in tissue macrophages under the pathologic circumstance of atheroma formation by Tangirala *et al.*, 36 *J. Lipid Res.* 2320-2328 (1995). The entire construct containing cathepsin S/K and the promoter is removed from the cloning vector and the purified DNA used to inject four-cell stage oocytes (C57 b/6) (20 µg per injection). The founder mice and the following progenies are characterized with tail DNA southern blot analysis with cathepsin cDNA probes. Crossbreeding of transgenic cathepsin S/K mice with LDLR^{-/-} mice is carried out in an animal facility. The homozygosity for cathepsin S/K transgene and LDLR^{-/-} alleles is tested with tail DNA southern blot analysis.

A strong promoter is required for cathepsin overexpression. We have found that mutations at the 3'-end of the original 4.5 kb mouse scavenger promoter can increase the reporter gene expression 100-1000-fold.

5 Coexpression and overexpression of both cathepsins in this transgenic mouse are verified by Southern and Western blot analyses and crossbreeding with LDLR^{-/-} mice. Coexpression and overexpression of both cathepsins in this transgenic line will induce atherogenesis, and the inhibitory and protective effect of cystatin C administration can be demonstrated by immunohistochemical comparison, as discussed above in EXAMPLE 1.

10 Alternatively, acute vascular injury may be created by placement of a nonconstrictive cuff around the mouse femoral artery, which results in smooth muscle cells migration, proliferation, and neointima formation. Briefly, mice are anesthetized on day 0 with nembutal and ketamine (0.04 mg/g for each). Under sterile conditions the femoral artery is dissected free and loosely sheathed with a nonocclusive, flexible cuff made of a polyethylene tube (3.0 mm long PE-90, 0.86 mm inner diameter and 1.27 mm
15 outer diameter, Becton Dickinson & Co.) which may then be ligated with a 5-0 silk thread. The wound is irrigated with saline, incision closed with running 3.0 silk suture, and topical antibiotic ointment applied. After recovery from anesthesia, the animals are given a standard diet and water *ad libitum*. Post-operative analgesia is administered using butorphenone (0.5-5.0 mg/kg s.c.). At day 14 after cuff placement, the animals are
20 re-anesthetized with ether, the chest cavity opened, and then sacrificed by placement of a 22-gauge butterfly angiocatheter into the left ventricle and in situ constant pressure (100 mm Hg) perfusion with 0.9% saline for 1 min. followed by fixation with 4% paraformaldehyde in 0.1 M phosphate buffer, pH 7.3, for 10 min. The right atrium may be opened laterally and functioned as the outflow tract. Both left and right femoral
25 arteries are then harvested, embedded in paraffin, and cut into transverse sections for routine light microscopy and immunohistochemistry. The inhibition of cathepsin S and cathepsin K at the site of vascular injury by systemic administration of cystatin C or

TGF- β_1 may then be examined. Cathepsin K activity, for example, may be examined by monitoring specific cathepsin K cleavage sites in native collagen using differential staining with monoclonal antibodies to the amino-terminal and distal helical regions of collagen (Johnatty *et al.*, 158 J. Immunol. 2327-333 (1997)).

5 Alternatively, acute vascular injury may be induced by carotid artery dilatation and endothelial cell denudation. Briefly, mice (C57/bl6, 8 weeks old weighing approximately 25 g) are anesthetized on day 0 by i.p. injection of ketamine (80 mg/kg) and xylazine (5 mg/kg). The skin overlying the right carotid artery is prepared for surgery and surgery performed using sterile techniques. The biological response to vascular injury may be investigated using an air-drying model. The animals are put on a standard diet and water *ad libitum* after surgery.

10 and post-operative analgesia is administered using butorphenol (0.5-5.0 mg/kg). At day 3, 7, or 28 the animals are re-anesthetized with ketamine and xylazine, the chest cavity opened, and the animal sacrificed by placement of a 22-gauge butterfly angiocatheter into the left ventricle for *in situ* constant pressure (100 mg) perfusion with 0.9% saline for 1 minute followed by fixation with 4% formaldehyde in 0.1M phosphate buffer, pH 7.3, for ten minutes. The right atrium may be opened laterally and functioned as the outflow tract. Both left and right carotid arteries are harvested, and embedded in paraffin for examination as discussed above in EXAMPLE 1.

15 Briefly, i.p. injections of cystatin C may be carried out every 48 hours. Spleen cells may be isolated and labeled with [125 I]-Z-Tyr-Ala-CHN₂, a cysteine protease active site probe, as previously described, and inhibition of cathepsins S and K monitored. Both cystatin C and TGF- β_1 compositions may be prepared as described above.

20 The foregoing description has been presented only for the purposes of illustration and is not intended to limit the invention to the precise form disclosed, but by the claims appended hereto.